INTERNATIONAL JOURNAL OF PHARMACEUTICS

VOL. 100 NO. 1–3 CONTENTS 8 NOVEMBER 1993

Cited in: Current Contents (Life Sciences; Clinical Practice); International Pharmaceutical Abstracts; Biological Abstracts; Chemical Abstracts; Excerpta Medica EMBASE; PASCAL M; Science Citation Index; Current Awareness in Biological Sciences

Announcement from the Publisher v

Editorial vii

Research Papers

The prediction of percutaneous absorption: I. Influence of the dermis on in vitro permeation models
O. Díez-Sales, E. Pérez-Sayas, A. Martín-Villodre and M. Herráez-Domínguez (Spain) 1

Effects of polymerization variables on PLGA properties: molecular weight, composition and chain structure
Mª.J. Dorta, O. Munguía and M. Llabrés (Spain) 9

Photochemical stability of biologically active compounds: V. Photochemical degradation of primaquine in an aqueous medium
S. Kristensen, A.-L. Grislingaas, J.V. Greenhill, T. Skjetne, J. Karlsen and H.H. Tønnesen (Norway, USA) 15

Influence of β-cyclodextrin concentration and polyacrylic acid molecular weight on swelling and release characteristics of metoclopramide-containing hydrogels
N. García-González, I.W. Kellaway, H. Blanco-Fuente, S. Anguiano-Igea, B. Delgado-Charro, F.J. Otero-Espinar and J. Blanco-Méndez (Spain, UK) 25

Mixture design of theophylline retard formulation
A. Kristl, D. Vojnovic, R. Karba, A. Mrhar, F. Rubessa and B. Zupaňi (Slovenia, Italy) 33

Vehicle effects on in vitro skin permeation of and stratum corneum affinity for model drugs caffeine and testosterone
F.P. Bonina, V. Carelli, G. Di Colo, L. Montenegro and E. Nannipieri (Italy) 41

Porous cellulose matrices – a novel excipient for the formulation of solid dosage forms
R. Davidson, H. Nyqvist and G. Ragnarsson (Sweden) 49

Study of the mechanisms of formation of nanoparticles and nanocapsules of polyisobutyl-2-cyanoacrylate
M. Gallardo, G. Couarraze, B. Denizot, L. Treupel, P. Couvreur and F. Puisieux (France) 55

Design and evaluation of buccoadhesive metoclopramide hydrogels composed of poly(acrylic acid) crosslinked with sucrose
N. García-González, I.W. Kellaway, H. Blanco-Fuente, S. Anguiano-Igea, B. Delgado-Charro, F.J. Otero-Espinar and J. Blanco-Méndez (Spain, UK) 65

Response surface optimization of high dose pellets by extrusion and spheronization
G.A. Hileman, S.R. Goskonda, A.J. Spalitto and S.M. Upadrashta (USA) 71

Gastrointestinal transit of pellets of differing size and density
G.M. Clarke, J.M. Newton and M.D. Short (UK) 81

Curcumin inhibits iron-dependent lipid peroxidation
Sreejayan and M.N.A. Rao (India) 93

Naproxen 1-alkylazacycloalkan-2-one esters as dermal prodrugs: in vitro evaluation
F.P. Bonina, L. Montenegro and F. Guerrera (Italy) 99

Selective coating of cylindrical matrices with a central hole. I. An interpretation of the swelling process
M.A. Vandelli and R. Cameroni (Italy) 107

Selective coating of cylindrical matrices with a central hole. II. An interpretation of the release process
M.A. Vandelli, G. Coppi and R. Cameroni (Italy) 115

Aerosolized aqueous suspensions of poly(L-lactic acid) microspheres
L.E. Masinde and A.J. Hickey (USA) 123

Evaluation of sustained-release granules of chlorphenesin carbamate in dogs and humans
M. Akimoto, A. Furuya, T. Maki, K. Yamada, T. Suwa and H. Ogata (Japan) 133

The influence of substitution type on the performance of methylcellulose and hydroxypropylmethycellulose in gels and matrices
K. Mitchell, J.L. Ford, D.J. Armstrong, P.N.C. Elliott, J.E. Hogan and C. Rostron (UK) 143

(Contents continued on p. 322)

0378-5173(19931108)100:1/3;1-E

(Contents continued from back cover)

The influence of concentration on the release of drugs from gels and matrices containing Methocel®
K. Mitchell, J.L. Ford, D.J. Armstrong, P.N.C. Elliott, C. Rostron and J.E. Hogan (UK) 155
The influence of drugs on the properties of gels and swelling characteristics of matrices containing methylcellulose or hydroxypropylmethylcellulose
K. Mitchell, J.L. Ford, D.J. Armstrong, P.N.C. Elliott, J.E. Hogan and C. Rostron (UK) 165
The influence of the particle size of hydroxypropylmethylcellulose K15M on its hydration and performance in matrix tablets
K. Mitchell, J.L. Ford, D.J. Armstrong, P.N.C. Elliott, J.E. Hogan and C. Rostron (UK) 175
A systematic study on the chemical stability of the novel indoloquinone antitumour agent EO9
J.D. De Vries, J. Winkelhorst, W.J.M. Underberg, R.E.C. Henrar and J.H. Beijnen (The Netherlands) 181
The effect of each of three physicochemical variables on solute uptake by polyvinylchloride infusion bags
P. Prayurnprohm and A.E. Polack (Australia) 189
Assessment of polysaccharide gels as drug delivery vehicles
L.E. McTaggart and G.W. Halbert (UK) 199
Interlot variability of carbomer 934
B. Pérez-Marcos, R. Martínez-Pacheco, J.L. Gómez-Amoza, C. Souto, A. Concheiro and R.C. Rowe (Spain, UK) 207
Evaluation of analytical methods: Liquid chromatography of cefalexin
C. Hendrix, Z. Yongxin, C. Van houtven, J. Thomas, E. Roets and J. Hoogmartens (Belgium) 213
Preparation and evaluation in vitro of solutions and o/w microemulsions containing levobunolol as ion-pair
M. Gallarate, M.R. Gasco, M. Trotta, P. Chetoni and M.F. Saettone (Italy) 219
Simultaneous quantitative determination of salbutamol sulfate and bromhexine hydrochloride in drug preparations by difference spectrophotometry
P. Parimoo, P. Umapathi and K. Ilango (India) 227
Transport and degradation characteristics of methotrexate dialkyl ester prodrugs across tape-stripped hairless mouse skin
J.J. Fort, Z. Shao and A.K. Mitra (USA) 233
Properties of hydroxypropylmethylcellulose granules produced by water spraying
C.-H. Liu, S.-C. Chen, Y.-H. Kao, C.-C. Kao, T.D. Sokoloski and M.-T. Sheu (Taiwan, ROC, USA) 241

Notes
The effect of magnesium trisilicate on proguanil absorption
C.O. Onyeji and C.P. Babalola (Nigeria) 249
Nil effect of cigarette smoking on ranitidine pharmacokinetics
Abd.E.M. Gharaibeh, H.N. Alkaysi, M.S. Salem and K.I.A. Gharaibeh (Jordan) 253
The effect of the physicochemical properties of a drug on its release from chitosonium malate matrix tablets
J. Akbuğa (Turkey) 257
Structure and behaviour of hydrophilic matrix sustained release dosage forms: 1. The origin and mechanism of formation of gas bubbles in the hydrated surface layer
C.D. Melia, A.R. Rajabi-Siahboomi, A.C. Hodsdon, J. Adler and J.R. Mitchell (UK) 263
Biological effects of a dipeptide methyl ester on *Echinococcus multilocularis* metacestodes in vivo
N. Walchshofer, M.-E. Sarciron, C. Arsac, S. Walbaum, J. Paris and A.-F. Petavy (France) 271
Effects on splenic, hepatic, hematological, and growth parameters following high-dose poloxamer 407 administration to rats
T.P. Johnston, H. Beris, Z.G. Wout and J.L. Kennedy (USA) 279

Author Index 285

Noticeboard 289

Master Index Vols. 91–100 291